

TABLE:PERCENTAGE OF WOMEN EXPERIENCING AN UNINTENDED PREGNANCY DURING THE FIRST YEAR OF A CONTRACEPTIVE METHOD

| Method | Perfect Use | Average Use |
|--|-------------|-------------|
| Levonorgestrel implants | 0.05 | 0.05 |
| Male sterilization | 0.10 | 0.15 |
| Female sterilization | 0.50 | 0.50 |
| Depo-Provera® (injectable progestogen) | 0.30 | 0.30 |
| Oral contraceptives | | 5 |
| Combined | 0.10 | NA |
| Progestin only | 0.50 | NA |
| IUD | | |
| Progesterone | 1.50 | 2.00 |
| Copper T 380A | 0.60 | 0.80 |
| Condom (male) without spermicide | 3 | 14 |
| (female) without spermicide | 5 | 21 |
| Cervical cap | | |
| Never given birth | 9 | 20 |
| Given birth | 26 | 40 |
| Vaginal Sponge | | |
| Never given birth | 9 | 20 |
| Given birth | 20 | 40 |
| Diaphragm with spermicidal cream or jelly | 6 | 20 |
| Spermicides alone (foam, creams, jellies, and vaginal suppositories) | 6 | 26 |
| Periodic abstinence (all methods) | 1-9* | 25 |
| Withdrawal | 4 | 19 |
| No contraception (planned pregnancy) | 85 | 85 |

NA -not available

*Depending on method (calender, ovulation symptothermal, post-ovulation

Adapted from Hatcher RA et al, *Contraceptive Technology: 17th Revised Edition*. NY,NY:

Ardent Medi, Inc, 1998